

# Get These Out of Your Family's Kitchen!

HYDROGENATED FATS OR OILS or PARTIALLY HYDROGENATED FATS OR OILS – Do not eat these.

If the label contains the words HYDROGENATED or PARTIALLY HYDROGENATED:

- do not eat it yourself.
- do not let your family or friends eat it.
- get it out of your family's kitchen.

HYDROGENATED or PARTIALLY HYDROGENATED FATS OR OILS – Read labels and avoid these substances like they are poisons...which they are.

## **“POISONS” IN OUR DIETS**

Why not consume partially hydrogenated fats or oils? Because by the definition of the “poison”, partially hydrogenated fats and oils are poisons.

Dorland's Medical Dictionary defines “Poison” as “any substance which, when relatively small amounts are ingested...has chemical action that may cause damage to structure or disturbance of functions, production symptomatology, illness, or death”.

Partially hydrogenated oils do not exist in nature. Partially hydrogenated fats and oils are processed versions of naturally occurring fats and oils. In nature, most dietary fats and oils exist in a structural form which is called the “cis” form. When these natural, cis form fats are processed by bubbling hydrogen gas through them at high temperatures, they become partially hydrogenated which changes their structure to the “trans” form. The natural cis fat has a bend and the processed trans fat is a straight molecule. The terms partially hydrogenated fat and trans fat will be used interchangeably. This difference in cis and trans shapes is of major significance. When eaten, fats and oils are incorporated into your cell membranes. Trans fats alter the configuration of these delicate structures. Further, trans fats interact with normal fat metabolism disturbing function in a most destructive manner. Hence, these substances meet the definition of a “poison”.

Trans fats interfere with important, normal functions by inhibiting enzymes which are necessary for your body's normal metabolism of fats and they keep doing it for a long time. When you eat normal cis fats, your body metabolizes half of them in 18 days. When you eat trans fats your body requires 51 days to metabolize half of them. This means that half of the trans fats you eat today will still be inhibiting essential enzyme systems in your body 51 days from now.

## **A MAJOR SOURCE OF CHRONIC DISEASE**

Many essential functions in our bodies depend on a group of hormones called prostaglandins (PGs) which are produced from fats in our diets. In a general sense there are good PGs and bad PGs. In our society most of the ill effects of chronic disease are promoted by or aggravated by the bad PGs called the PG 2 family.

This includes:

- **Heart attacks & Strokes**
- **Cardiovascular disease**
- **Cancer**
- **Inflammatory conditions & Autoimmune diseases**

The PG family is derived directly from naturally occurring fat found in red meat, shellfish, and dairy products. PG family reactions include: increasing blood clotting, increasing blood pressure, increasing cholesterol and causing other heart attack risk factors to be made worse. PG 2 family substances increase inflammatory activity which leads to tissue destruction associated with

everything from trauma to autoimmune diseases. PG 2 family chemicals also cause a decrease in natural killer cells which are necessary for your body to fight off cancer. Because of this facts, tumors increase in size under the influence of PG 2.

The good PG families counteract the bad effects of the PG 2 family. They decrease blood clotting, decrease blood pressure, decrease cholesterol, decrease inflammation, and increase natural killer cell activity which is necessary to fight tumors.

Trans fats block good PGs production, and by default, the bad PG 2 substances are produced unopposed. The PG 2 imbalance created by the consumption of partially hydrogenated fats contributes to the production of chronic disease which is the enigma of our modern society.

There are dozens of references in scientific literature documenting the ill effects of trans fatty acids especially in relation to heart disease. If you have difficulty in convincing someone of the value in avoiding trans fats, some of the extensive references now available in the scientific literature are listed at the end of this pamphlet.

### **DAILY, NAGGING SYMPTOMS**

There are also other symptoms which are created by the bad PGs when trans fats inhibit the balancing effects of the good PGs. These include:

- **Headaches**
- **Joint pain & Back pain**
- **Arthritis**
- **Asthma**
- **Skins Problems**
- **Hot Flashes**
- **Premenstrual syndrome & Menstrual cramps**

These are just a few of the symptoms which are related to PG 2 imbalances created by eating partially hydrogenated fats and oils.

Millions of people take aspirin, acetaminophen, and other non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen or naproxen to alleviate these symptoms. These drugs work by blocking PG production. Avoidance of partially hydrogenated trans fats and consumption of natural fats and oils can often turn around these common, nagging symptoms in as little as 3 to 4 weeks without any other intervention.

It has been said that, in the history of the world, there has never been an aspirin deficiency or an NSAID deficiency. But there are millions of people who regularly consume partially hydrogenated fats and oils who suffer ill effects and then mask the symptoms with these drugs. *If you, your family, or your friends achieve symptom relief with aspirin, acetaminophen, or any other NSAID, it is almost certain that there is a fatty acid imbalance in the body causing the symptoms in the first place.*

Nutritional supplementation with the essential fatty acids can be helpful in reducing symptoms. This is especially true if symptom relief is achieved by aspirin or similar substances. The good PGs can be supplemented with black currant seed oil, evening primrose oil, borage oil, flaxseed (linseed) oil, and/or fish oils such as those containing EPA (eicosapentaenoic acid). Ask your doctor about which of these may be useful to you.

### **RELIEF OF "NORMAL" HEADACHES**

Although supplementation is useful, return to normal function will not be seen until the partially hydrogenated fats and oils are eliminated from the diet. Many people put up with or

medicate for daily nagging symptoms which they wrongly assume are normal. There are no such thing as normal headaches, normal backaches, normal arthritis, normal menstrual cramps, etc.

All of these symptoms have been helped by total avoidance of partially hydrogenated fats and oils. You will be elated to learn that daily, nagging symptoms are not “normal”, nor due to “normal aging” or “just getting older”. Due to the prolonged life of trans fatty acids, you must be both diligent in avoiding them and patient for the changes to take place. Within a few weeks or a couple of months the results are usually noticeable and often quite gratifying.

### **READ LABELS AS IF YOUR LIFE DEPENDED ON IT**

Read labels. Read labels as if your life depended on it. It does. As people say when they first look to avoid these poisons, “these fats are in everything on the grocery store shelves!” Well, not quite everything. But partially hydrogenated oils are found in margarines, vegetable shortenings, mot chips, cookies, candy, cakes, popcorn and other similar snack foods, and are used in food preparation by most fat food companies. Also, many peanut butters, canned soups, and some so-called “health food” products are made with trans fats. Typical ingredients sections of labels include such names as “partially hydrogenated soybean oil”, “hydrogenated vegetable oil”, “partially hydrogenated cottonseed oil”, and so on.

Unfortunately there are some doctors who are unaware of the mass of recent research evidence indicating the trans fats as a source of increased risk of heart disease, much less the other adverse effects of consuming these poisonous substances. In fact, many doctors still recommend margarine instead of butter for patients trying to prevent or improve heart disease. The fact is that trans fats increase cardiac risk factors twice as much as saturated fat in the diet!

Still, many patients blindly follow their misinformed doctor’s advice and are unwittingly consuming foods thinking they will decrease their risk for heart disease. In fact, these foods will actually significantly worsens their cardiac risk factors, not to mention the other problems they create. We must stop the insanity of slowly poisoning ourselves and our society by replacing natural fats with processed trans fats.

Many of us, our family members, and our friends have nagging symptoms or major degenerative problems which we more or less take for granted. We refer to these symptoms as “normal” or “typical”, which they are not. When they recur, we feel victims of an imperfect world, or we write them off to the aging process. They are tangible reasons for most everything we, our families, and our friends experience. The ingestion of hydrogenated and partially hydrogenated fats and oils contributes to the common aches and pains of daily life as well as progressive, degenerative processes and life ending illnesses. These symptoms can be changed and the quality of life improved by simply avoiding hydrogenated and partially hydrogenated fats and oils.

Now you can see why we say “Get these out of your family’s kitchen!”

**FRIENDS DON'T LET FRIENDS EAT PARTIALLY HYDROGENATED FATS!**

## REFERENCES

1. Danneneberg AJ, Redienberg MM. Clinical pharmacological therapeutics, 1994;55:5-9
2. McGiff JC et al. Hypertension 1991; (Suppl.3):18(sup III):III-150-III-157
3. Saloman P, et al. J clin gastroenterology, 1990;12(2):157-161
4. Sperling, RI. Nutrition in rheumatic diseases/rheumatic disease clinics of North America/nutrition and rheumatic diseases, May 1991;17(2):373-389
5. Callejari PE and Surier, RB. Nutrition and rheumatic diseases/rheumatic disease clinics of North America, May 1991;17(2):415-425
6. Narisawa, T; et al. Diseases of colon and rectum, Oct 1990;33(10):840-845
7. El-Hakm IK, et al. Brit j maxillofacial surg. 1990;29:155-159
8. Axelrod L; et al. Diabetes care, Jan 1994;17(1):37
9. Zhu B; et al. the American Heart Journal. 04/94;127(No.4/part 1):744-755
10. Goodnight, SH. Archives of pathology and laboratory medicine, Jan 1993;117:102-105
11. Engler MB. J cardiovasc nurs, 1994;8(3):53-67
12. Nordoy A; et al. Am j clin nutr, 1993;57:634-639
13. Parkinson A. Am j clin nutr, 1994;59:384-388
14. Brzeski, M et al. British journal of rheumatology, 1991;30:370-372
15. Ascherio A; et al. Circulation, 1994;89:94-101
16. Willett WC; et al. Lancet, 03/06/93;341:581-585
17. Hurley J, Schmidt S. Nutrition action, May. 1994;1,9
18. Litin L, Sacks F. New Engl j med, Dec 23, 1993;329(26):1969-1970